

Analysis of genome-wide genotype data

Acknowledgement: Several slides based on a lecture course given by Jonathan Marchini & Chris Spencer, Cape Town 2007

Introduction & definitions

- Allele: A version of a genetic locus
- Single Nucleotide Polymorphism (SNP): Common variants (>1% frequency of minor allele?)
- An allele can be the
 - Minor allele (the less frequent)
 - Ancestral
 - Derived (opposite to ancestral)
- At a polymorphic locus an individual can be
 - Homozygous (two alleles identical)
 - Heterozygous

Human variation


A pair of randomly sampled human chromosomes differ at about 1 site per 1,200

A total of ~ 10 million common SNPs in human (account for 90% of human variation)

Many variants account for the rest

Linkage, haplotypes and HapMap

Haplotype: A set of alleles co-occurring on a chromosome (contraction of haploid genotype)



Linkage, haplotypes and HapMap


Haplotype: A set of alleles co-occurring on a chromosome (contraction of haploid genotype)

Alleles located close to one another are inherited together (said to be linked or in linkage disequilibrium)

This pattern is disrupted by recombination (shuffles chromosomes)


But recombination finite and time since origin of humans (~150,000 years) insufficient to break down all linkage => haplotype blocks

Recombination breaks down LD



- Recombination breaks down LD by making mosaic of existing haplotypes.

LD and association mapping -> Genome-wide association studies (GWAS)




Disease mutation
arises on single
haplotype

Increase in
frequency by drift.


Shuffling occurs
due to
recombination

Why? Hotspots of recombination



- Experimental evidence shows that recombination is clustered in hotspots
- Hotspots are interspersed with long cold spots


Recombination and LD



HapMap Consortium 2005


- Recombination hotspots demark break down in LD
- Inter-hotspot expanses have multiple markers in strong LD

Implications of recombination hotspots



- Block-like structure to LD across the human genome
- Few markers may tag a large number of SNPs
- More markers needed in recombination hotspots

Relationship between haplotype blocks and recombination hotspots



McVean, Spencer Chaix, 2005

Imputation and Tag SNPs

Because of linkage the allele at a given SNP can predict the allele at a linked SNP

Therefore sufficient to genotype a subset of SNPs (Tag SNPs) and predict the rest

The HapMap project (begun in 2002)

Aims

Map variation across human populations

Determine tag SNPs (~600,000)

- one of the major reasons for the HapMap project
- several methods have been proposed to choose the optimal set of tags

Which leads to reduced cost of genotyping individuals

Facilitates research into genetic disorders

Make all the data publicly available

Haplotype blocks

Entrez genes


NM_015158

ANKRD15: ankyrin repeat domain protein 15 isoform a


NM_153186

ANKRD15: ankyrin repeat domain protein 15 isoform b

CEU Phased Haplotypes



YRI Phased Haplotypes



Haplotype data from HapMap


[illegible]

Generation of 'genome-wide' genotype data


Generally carried out using 'microarray' technologies

- small glass slides typically containing millions of 'spots' consisting of immobilized nucleic acid sequences (or probes)
- genomic DNA is first fragmented (using enzymes) and labelled with a fluorescent dye
- genomic DNA washed over slide, hybridizes to complementary probes
- unhybridized DNA washed off
- fluorescent signal detected by scanning with a laser

Microarray




Genotype Calling



E.g. Chiamo (Marchini et al.)

Model



Example: Wellcome Trust Case Control Consortium, *Nature* 2007

Investigated genetics of seven common complex disorders

~ 17,000 samples, ~500,000 SNPs

Used a logistic regression model. Let Z_i be the genotype of individual i , encoded as 0, 1 or 2 (depending on number of copies of the non-reference allele)

$$M_0 : \log\left(\frac{p}{1-p}\right) = \mu$$


$$M_1 : \log\left(\frac{p}{1-p}\right) = \mu + \gamma Z_i$$

$$M_2 : \log\left(\frac{p}{1-p}\right) = \mu + \gamma I(Z_i = 1) + \phi(2\gamma I(Z_i = 2))$$


We can calculate Bayes factors, as $\frac{P(D|M_1)}{P(D|M_0)}$

$$= \frac{\int P(D|\theta_1, M_1)P(\theta_1|M_1)d\theta_1}{\int P(D|\theta_0, M_0)P(\theta_0|M_0)d\theta_0}$$


e.g. Results of Wellcome Trust Case Control Consortium (Nature, 2007)



Accounting for multiple testing a key issue in GWAS



T2D hit region, chromosome 10



WTCCC