

Proof of the fundamental Theorem of Calculus

We want to prove

$$\int_S dw = \int_{\partial S} w$$

where w is a 0-form.

For simplicity let's consider the case of $n=2$ variables.

Proof

Choose $w = F(x, y)$.

Suppose $x = g(t)$, $y = h(t)$ is some parameterization of S as variable t varies from t_0 to t_1 .



$$\int_S dw \approx \int_S F_x(x, y) dx + F_y(x, y) dy$$

$$= \int_{t_0}^{t_1} F_x(g(t), h(t)) g'(t) dt + F_y(g(t), h(t)) h'(t) dt$$

$$= \int_{t_0}^{t_1} (F_x(g(t), h(t)) g'(t) + F_y(g(t), h(t)) h'(t)) dt$$

chain rule

$$= \int_{t_0}^{t_1} \left(\frac{dF}{dt} \right) dt$$

$$= F(g(t_1), h(t_1)) - F(g(t_0), h(t_0))$$

FTC in
one
variable

$$= F(Q) - F(P)$$

$$= \int_S w$$

QED

Summary of 1-forms and an introduction to 2-forms

- A 1-form is an expression such as

$$\omega = A dx + B dy$$

that can be integrated over oriented curves. (= oriented 1-dimensional regions)

- A 2-form is an expression such as

$$\omega = A dx \wedge dy + B dy \wedge dz + C dz \wedge dx$$


that can be "integrated" over 2-dimensional "oriented regions".

• Integrals of 1-forms are just ^{limits} of sums of integrals of constant 1-forms over oriented straight line segments.



• Integrals of 2-forms are just limits of sums of "constant 2-forms" over "oriented planar triangles".

Oriented Planar Triangles

Three points on a plane determine a triangle.



An orientation of a triangle is specified by an arrow



corresponding to one of two possible directions of rotation.

The positive side of an oriented triangle is the one

is the one for which the arrow denotes anti-clockwise rotation,

An orientation is just an ordering of the vertices. The ordering PQR denotes the second triangle above, so too does RQP , so too does QPR .
