

sol

$$u_x = 2 + e^x \sin y$$

$$u_{xx} = e^x \sin y$$

$$u_y = 1 + e^x \cos y$$

$$u_{yy} = -e^x \sin y$$

$$u_{xx} + u_{yy} = 0 \quad \checkmark$$

lec 8

- If two harmonic functions

$u(x, y)$ and $v(x, y)$

satisfy the C-R equations

in a domain D ,

they are the real and

imaginary parts of an

analytic function f in D .

$$[f(z) = u(x, y) + i v(x, y)]$$

- $v(x, y)$ is said to be the conjugate harmonic function of u in D

(This has nothing to do with the "conjugate" \bar{z})

Ex Earlier we showed

$$u(x, y) = 2x + y + e^x \sin(y)$$

is harmonic.

Find a conjugate harmonic function v of u .

Sol

$$\frac{\partial u}{\partial x} = 2 + e^x \sin y$$

As $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$

we can say: $\frac{\partial v}{\partial y} = 2 + e^x \sin y$

or $v(x, y) = 2y - e^x \cos y + h(x)$ *

$$\frac{\partial u}{\partial y} = 1 + e^x \cos y$$

As $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$

we can say $\frac{\partial v}{\partial x} = -1 - e^x \cos y$

$$\text{or } v(x, y) = -x - e^x \cos y + 2y$$

$$\text{Rewriting } * v(x, y) = h(x) - e^x \cos y + 2y$$

$$\text{Therefore, } h(x) = -x$$

$$\text{and } l(y) = 2y$$

$$\text{So } v(x, y) = -x - e^x \cos y + 2y$$

is a conjugate harmonic function of $u(x, y)$.

Note: we could have picked

$$h(x) = -x + c_1$$

or

$$l(y) = 2y + c_2$$

$$c_1, c_2 \in \mathbb{R}$$

So, there are infinitely many conjugate harmonic functions of $u(x, y)$, given by

$$v(x, y) = -x - e^x \cos y + 2y + c_2$$