


Recall

K a simplicial complex

Example



$$\begin{array}{l} 4 \mid V = \{0, 1, 2, 3, 4\} \\ K = \{ \{0\}, \{1\}, \{2\}, \\ \quad \{3\}, \{4\}, \\ \quad \{0, 1\}, \{0, 2\} \\ \quad \{0, 3\}, \{1, 2\} \\ \quad \{2, 3\}, \\ \quad \{0, 1, 2\} \} \end{array}$$

$C_n K$ a vector space (over say \mathbb{R})
with one basis element

e_σ
for each n -simplex

$$\sigma = \{v_0, v_1, \dots, v_n\} \in K.$$

$d_n: C_n K \rightarrow C_{n-1} K$ is a linear homⁿ
given by

$$d_n(e_\sigma) = \sum_{i=0}^n (-1)^i e_{\sigma \setminus \{v_i\}}$$

Defn The degree n Betti number
of K is

$$B_n = \dim(K_n) - \dim(\text{image } d_n)$$

β_n measures, in some sense, 'the number of n -dimensional holes' in K .

β_n measures, in some sense, 'the number of n -dimensional holes' in K .

A '0-dimensional hole' in K is defined to be a connected component of K .

β_n measures, in some sense, 'the number of n -dimensional holes' in K .

A '0-dimensional hole' in K is defined to be a connected component of K .
One can prove that β_0 equals the number of connected components of K .

β_n measures, in some sense, 'the number of n -dimensional holes' in K .

A '0-dimensional hole' in K is defined to be a connected component of K .
One can prove that β_0 equals the number of connected components of K .

The clique complex χK_ϵ depends on $\epsilon \geq 0$.

β_n measures, in some sense, 'the number of n -dimensional holes' in K .

A '0-dimensional hole' in K is defined to be a connected component of K . One can prove that β_0 equals the number of connected components of K .

The clique complex K_ϵ depends on $\epsilon \geq 0$. We'd like to consider all $0 \leq \epsilon \leq \infty$ and determine the number of n -dimensional holes in K_ϵ that persist over a long range of values of ϵ .

β_n measures, in some sense, 'the number of n -dimensional holes' in K .

A '0-dimensional hole' in K is defined to be a connected component of K . One can prove that β_0 equals the number of connected components of K .

The clique complex K_ϵ depends on $\epsilon \geq 0$. We'd like to consider all $0 \leq \epsilon \leq \infty$ and determine the number of n -dimensional holes in K_ϵ that persist over a long range of values of ϵ .

This idea will lead us to a definition of **persistent Betti numbers**.

β_n measures, in some sense, 'the number of n -dimensional holes' in K .

A '0-dimensional hole' in K is defined to be a connected component of K . One can prove that β_0 equals the number of connected components of K .

The clique complex K_ϵ depends on $\epsilon \geq 0$. We'd like to consider all $0 \leq \epsilon \leq \infty$ and determine the number of n -dimensional holes in K_ϵ that persist over a long range of values of ϵ .

This idea will lead us to a definition of **persistent Betti numbers**.

But let's first consider the idea informally, via a potential application, before studying the precise mathematical definition of persistent homology.

A general problem

Given a set S of points randomly sampled from an unknown manifold M , what can we infer about the topology of M ?

A general problem

Given a set S of points randomly sampled from an unknown manifold M , what can we infer about the topology of M ?

For instance, $S \subset M \subset \mathbb{E}^2$.



One approach to the problem

Repeatedly “thicken” the set S to produce a sequence of inclusions

$$S = S_1 \subset S_2 \subset S_3 \subset \dots$$

and then search for “persistent” topological features in the sequence.

One approach to the problem

Repeatedly “thicken” the set S to produce a sequence of inclusions

$$S = S_1 \subset S_2 \subset S_3 \subset \dots$$

and then search for “persistent” topological features in the sequence.



One approach to the problem

Repeatedly “thicken” the set S to produce a sequence of inclusions

$$S = S_1 \subset S_2 \subset S_3 \subset \dots$$

and then search for “persistent” topological features in the sequence.



One approach to the problem

Repeatedly “thicken” the set S to produce a sequence of inclusions

$$S = S_1 \subset S_2 \subset S_3 \subset \dots$$

and then search for “persistent” topological features in the sequence.



Betti numbers

$\beta_0(X)$ = number of path components of X

Betti numbers

$\beta_0(X)$ = number of path components of X

	S_1	S_2	S_3	S_4	S_5	S_6	S_7	S_8
β_0	478	32	9	2	1	1	1	1

Betti numbers

$\beta_0(X)$ = number of path components of X

$\beta_1(X)$ = number of "1-dimensional holes" in X

	S_1	S_2	S_3	S_4	S_5	S_6	S_7	S_8
β_0	478	32	9	2	1	1	1	1
β_1	0	115	18	4	1	1	1	1

Betti numbers




$\beta_0(X)$ = number of path components of X

$\beta_1(X)$ = number of "1-dimensional holes" in X




	S_1	S_2	S_3	S_4	S_5	S_6	S_7	S_8
β_0	478	32	9	2	1	1	1	1
β_1	0	115	18	4	1	1	1	1

These numbers are consistent with the sample coming from some region with the homotopy type of a circle.

During an inclusion $S_i \hookrightarrow S_j$ holes can






During an inclusion $S_i \hookrightarrow S_j$ holes can



β_n^{ij} = number of n -dimensional holes in S_i that persist to S_j

During an inclusion $S_i \hookrightarrow S_j$ holes can



β_n^{ij} = number of n -dimensional holes in S_i that persist to S_j

β_0^{ij} = number of connected components in S_i that persist to S_j

Bar codes


The matrix (β_n^{ij}) can be represented by a graph with horizontal edges and vertices arranged in columns.

The i th column has $\beta_n^{ii} = \beta_n(S_i)$ vertices.

There are β_n^{ij} paths from the i th column to the j th column.




β_1 bar code for
our example



β_0 bar code for our example
(first column cropped)

Example Consider a sample S of 750 points selected at random from two "quarter segments" of a torus.



$$S \subset \text{Torus} \subset \mathbb{R}^3$$

Defn The degree n Betti number
of K is

$$\beta_n = \dim(Ker d_n) - \dim(\text{image } d_{n+1})$$

Defn The degree n Betti number of K is

$$\beta_n = \dim(K_n) - \dim(\text{image } d_n)$$

It is convenient to view β_n as the dimension of a certain vector space.

Lemma

$$d_n(d_{n-1}e_\sigma) = 0$$

for all n -simplices σ

proof exercise

Lemma

$$d_n(d_{n+1}e_\sigma) = 0$$

for all $n+1$ -simplices σ

Proof exercise

Thus

$$\text{Im } d_{n+1} \subseteq \ker d_n$$

Defn The degree n homology
of K is the vector space

$$H_n(K) = \frac{\ker d_n}{\operatorname{Im} d_{n+1}}.$$

Defn The degree n homology
of K is the vector space

$$H_n(K) = \frac{\ker d_n}{\operatorname{im} d_{n+1}}.$$

From this definition we see

$$B_n = \dim(H_n(K))$$

observation: If L is a subcomplex of the simplicial complex K (i.e. L is a simplicial complex with $L \subset K$) then

$C_n L$ is a sub vector space of $C_n K$.

In fact we have a diagram

$$\begin{array}{ccc}
 \vdots & & \vdots \\
 \downarrow & & \downarrow \\
 C_3 L & \longrightarrow & C_3 K \\
 \downarrow d_3 & & \downarrow d_3 \\
 C_2 L & \longrightarrow & C_2 K \\
 \downarrow d_2 & & \downarrow d_2 \\
 C_1 L & \longrightarrow & C_1 K \\
 \downarrow d_1 & & \downarrow d_1 \\
 C_0 L & \longrightarrow & C_0 K
 \end{array}$$

This diagram induces a homomorphism of vector spaces

$$H_n(L) \longrightarrow H_n(K)$$

which is not in general injective.