

Natural Images

Natural Images


van Hateren & van der Schaaf: 4167 digital photos of random outdoor scenes.

Natural Images


van Hateren & van der Schaaf: 4167 digital photos of random outdoor scenes.



Mumford, Lee, Pedersen: 8 000 000 random high-contrast 3×3 patches in \mathbb{E}^9 .



Mumford, Lee, Pedersen: 8 000 000 random high-contrast 3×3 patches in \mathbb{E}^9 .



By normalizing with respect to brightness and contrast the patches are projected onto a set of points \mathcal{M} in a topological seven-sphere $S^7 \subset \mathbb{E}^8$.

Carlsson, Ishikano, de Silva, Zomorodian: consider high-density subsets $\mathcal{M}(k) \subset \mathcal{M}$.

Carlsson, Ishikhanov, de Silva, Zomorodian: consider high-density subsets $\mathcal{M}(k) \subset \mathcal{M}$.

Codensity function: $\delta_k(x)$ is the Euclidean distance from $x \in \mathcal{M}$ to its k th nearest neighbour in \mathcal{M} .


Carlsson, Ishkhanov, de Silva, Zomorodian: consider high-density subsets $\mathcal{M}(k) \subset \mathcal{M}$.

Codensity function: $\delta_k(x)$ is the Euclidean distance from $x \in \mathcal{M}$ to its k th nearest neighbour in \mathcal{M} .

$\mathcal{M}(k)$ contains the 25% of patches with least $\delta_k(x)$.


The β_1 bar code for a random sampling of $\mathcal{M}(300)$ yields a single persistent homology class.

The β_1 bar code for a random sampling of $\mathcal{M}(300)$ yields a single persistent homology class.




The β_1 bar code for a random sampling of $\mathcal{M}(15)$ yields five persistent homology classes.



The β_1 bar code for a random sampling of $\mathcal{M}(15)$ yields five persistent homology classes.



The β_1 bar code for a random sampling of $\mathcal{M}(15)$ yields five persistent homology classes.





The β_1 bar code for a random sampling of $\mathcal{M}(15)$ yields five persistent homology classes.



The β_2 bar codes are less robust, but seem to indicate a single homology class with mod 2 coefficients.


The β_2 bar codes are less robust, but seem to indicate a single homology class with mod 2 coefficients.

$$H_2(\text{Klein bottle}, \mathbb{Z}_2) = \mathbb{Z}_2$$




The β_2 bar codes are less robust, but seem to indicate a single homology class with mod 2 coefficients.


$$H_2(\text{Klein bottle}, \mathbb{Z}_2) = \mathbb{Z}_2$$



The β_2 bar codes are less robust, but seem to indicate a single homology class with mod 2 coefficients.

$$H_2(\text{Klein bottle}, \mathbb{Z}_2) = \mathbb{Z}_2$$





Theorem. [Edelsbrunner, Chazal, ...] For two finite metric spaces S, S' and $n \geq 0$ we have

$$d_{BottleNeck}(\beta_n^{**}(S), \beta_n^{**}(S')) \leq d_{GromovHausdorff}(S, S') .$$

Let (X, d_X) and (Y, d_Y) be finite metric spaces.

Let (X, d_X) and (Y, d_Y) be finite metric spaces.

By a **correspondence** between them we mean a set $C \subset X \times Y$ such that both projections $C \rightarrow X$ and $C \rightarrow Y$ are surjective.

Let (X, d_X) and (Y, d_Y) be finite metric spaces.

By a **correspondence** between them we mean a set $C \subset X \times Y$ such that both projections $C \rightarrow X$ and $C \rightarrow Y$ are surjective.

Let $\Gamma(X, Y)$ denote the set of all correspondences between X and Y .

Let (X, d_X) and (Y, d_Y) be finite metric spaces.

By a **correspondence** between them we mean a set $C \subset X \times Y$ such that both projections $C \rightarrow X$ and $C \rightarrow Y$ are surjective.

Let $\Gamma(X, Y)$ denote the set of all correspondences between X and Y .

Define the **Gromov-Hausdorff distance** between X and Y to be

$$d_{GH}(X, Y) = \frac{1}{2} \inf_{C \in \Gamma(X, Y)} \sup_{(x, y), (x', y') \in C} |d_X(x, x') - d_Y(y, y')|.$$

A bounded closed interval $I = [a, b] \subset \mathbb{R}$ has length $b - a$.

A bounded closed interval $I = [a, b] \subset \mathbb{R}$ has **length** $b - a$.

For $\delta \geq 0$ define $\tau_\delta(I) = \{r \in \mathbb{R} : |r - s| \leq \delta \text{ for some } s \in I\}$.

A bounded closed interval $I = [a, b] \subset \mathbb{R}$ has **length** $b - a$.

For $\delta \geq 0$ define $\tau_\delta(I) = \{r \in \mathbb{R} : |r - s| \leq \delta \text{ for some } s \in I\}$.

A **barcode** is a finite multiset of bounded closed intervals in \mathbb{R} .

A bounded closed interval $I = [a, b] \subset \mathbb{R}$ has **length** $b - a$.

For $\delta \geq 0$ define $\tau_\delta(I) = \{r \in \mathbb{R} : |r - s| \leq \delta \text{ for some } s \in I\}$.

A **barcode** is a finite multiset of bounded closed intervals in \mathbb{R} .

For two barcodes C, D define a **matching** to be a bijection $C \supseteq S \leftrightarrow T \subseteq D$ between subsets S, T of C, D respectively.

A bounded closed interval $I = [a, b] \subset \mathbb{R}$ has **length** $b - a$.

For $\delta \geq 0$ define $\tau_\delta(I) = \{r \in \mathbb{R} : |r - s| \leq \delta \text{ for some } s \in I\}$.

A **barcode** is a finite multiset of bounded closed intervals in \mathbb{R} .

For two barcodes C, D define a **matching** to be a bijection $C \supseteq S \leftrightarrow T \subseteq D$ between subsets S, T of C, D respectively.

Let C^δ denote the multiset of those intervals in C of length at least δ .

A bounded closed interval $I = [a, b] \subset \mathbb{R}$ has **length** $b - a$.

For $\delta \geq 0$ define $\tau_\delta(I) = \{r \in \mathbb{R} : |r - s| \leq \delta \text{ for some } s \in I\}$.

A **barcode** is a finite multiset of bounded closed intervals in \mathbb{R} .

For two barcodes C, D define a **matching** to be a bijection $C \supseteq S \leftrightarrow T \subseteq D$ between subsets S, T of C, D respectively.

Let C^δ denote the multiset of those intervals in C of length at least δ .

Define a δ -**matching** between barcodes C and D to be a bijection $C^{2\delta} \leftrightarrow D^{2\delta}$ such that: the bijection associates each interval in $I \in C^{2\delta}$ with an interval in $\sigma(I) \in D^{2\delta}$; if $\sigma(I) = J$ then $J \subset \tau_\delta(I)$ and $I \subset \tau_\delta(J)$.

A bounded closed interval $I = [a, b] \subset \mathbb{R}$ has **length** $b - a$.

For $\delta \geq 0$ define $\tau_\delta(I) = \{r \in \mathbb{R} : |r - s| \leq \delta \text{ for some } s \in I\}$.

A **barcode** is a finite multiset of bounded closed intervals in \mathbb{R} .

For two barcodes C, D define a **matching** to be a bijection $C \supseteq S \leftrightarrow T \subseteq D$ between subsets S, T of C, D respectively.

Let C^δ denote the multiset of those intervals in C of length at least δ .

Define a δ -**matching** between barcodes C and D to be a bijection $C^{2\delta} \leftrightarrow D^{2\delta}$ such that: the bijection associates each interval in $I \in C^{2\delta}$ with an interval in $\sigma(I) \in D^{2\delta}$; if $\sigma(I) = J$ then $J \subset \tau_\delta(I)$ and $I \subset \tau_\delta(J)$.

Now define the **bottleneck distance** between two barcodes C, D to be

$$d_B(C, D) = \inf\{\delta : \text{there is a } \delta\text{-matching between } C \text{ and } D\}.$$