


Voronoi tessellation for set  $S \subset \mathbb{R}^2$  of 20 points.



By Balu Ertl - Own work,  
<https://commons.wikimedia.org/>


Consider  $S = \mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$ ,  $p = (0, 0) \in S$

Consider  $S = \mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$ ,  $p = (0, 0) \in S$

$$V(p) = \{v \in \mathbb{R}^2 : \|p - v\| < \|p' - v\| \text{ for all } v \in S\}$$


Consider  $S = \mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$ ,  $p = (0, 0) \in S$

$$V(p) = \{v \in \mathbb{R}^2 : \|p - v\| < \|p' - v\| \text{ for all } v \in S\}$$




Consider  $S = \mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$ ,  $p = (0, 0) \in S$

$$V(p) = \{v \in \mathbb{R}^2 : \|p - v\| < \|p' - v\| \text{ for all } v \in S\}$$




Consider  $S = \mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$ ,  $p = (0, 0) \in S$

$$V(p) = \{v \in \mathbb{R}^2 : \|p - v\| < \|p' - v\| \text{ for all } v \in S\}$$




Consider  $S = \mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$ ,  $p = (0, 0) \in S$

$$V(p) = \{v \in \mathbb{R}^2 : \|p - v\| < \|p' - v\| \text{ for all } v \in S\}$$



Consider  $S = \mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$ ,  $p = (0, 0) \in S$

$$V(p) = \{v \in \mathbb{R}^2 : \|p - v\| < \|p' - v\| \text{ for all } v \in S\}$$





Consider  $S = \mathbb{Z} \times \mathbb{Z} \subset \mathbb{R}^2$ ,  $p = (0, 0) \in S$

$$V(p) = \{v \in \mathbb{R}^2 : \|p - v\| < \|p' - v\| \text{ for all } v \in S\}$$

