

Computational Homology I

Berlin, August 2010

Graham Ellis
NUI Galway, Ireland

G = discrete group

X = any contractible space on which G acts freely

$$H_n(G, \mathbb{Z}) = H_n(X/G, \mathbb{Z})$$

G = discrete group

X = any contractible space on which G acts freely

$$H_n(G, \mathbb{Z}) = H_n(X/G, \mathbb{Z})$$

Definition is computational:

1. construct X and X/G efficiently
2. then compute $H_n(X/G, \mathbb{Z})$.

G = discrete group

X = any contractible space on which G acts freely

$$H_n(G, \mathbb{Z}) = H_n(X/G, \mathbb{Z})$$

Definition is computational:

1. construct X and X/G efficiently
2. then compute $H_n(X/G, \mathbb{Z})$.

Step 2 used in applied topology.

One problem from applied topology

Given a set S of points randomly sampled from an unknown manifold M , what can we infer about the topology of M ?

One problem from applied topology

Given a set S of points randomly sampled from an unknown manifold M , what can we infer about the topology of M ?

For instance, $S \subset M \subset \mathbb{E}^2$.



One approach to the problem

Repeatedly “thicken” the set S to produce a sequence of inclusions

$$S = S_1 \subset S_2 \subset S_3 \subset \cdots$$


and then search for “persistent” topological features in the sequence.

One approach to the problem

Repeatedly “thicken” the set S to produce a sequence of inclusions

$$S = S_1 \subset S_2 \subset S_3 \subset \cdots$$

and then search for “persistent” topological features in the sequence.




s1

One approach to the problem


Repeatedly “thicken” the set S to produce a sequence of inclusions

$$S = S_1 \subset S_2 \subset S_3 \subset \dots$$

and then search for “persistent” topological features in the sequence.



s1



s2

One approach to the problem

Repeatedly “thicken” the set S to produce a sequence of inclusions

$$S = S_1 \subset S_2 \subset S_3 \subset \dots$$

and then search for “persistent” topological features in the sequence.



s1



s2



s3



s4



s5



s6



s7



s8

Betti numbers

$\beta_0(X)$ = number of path components of X

Betti numbers

$\beta_0(X)$ = number of path components of X

	S_1	S_2	S_3	S_4	S_5	S_6	S_7	S_8
β_0	478	32	9	2	1	1	1	1

Betti numbers

$\beta_0(X)$ = number of path components of X

$\beta_1(X)$ = number of "1-dimensional holes" in X

	S_1	S_2	S_3	S_4	S_5	S_6	S_7	S_8
β_0	478	32	9	2	1	1	1	1
β_1	0	115	18	4	1	1	1	1

Betti numbers


$\beta_0(X)$ = number of path components of X

$\beta_1(X)$ = number of "1-dimensional holes" in X


	S_1	S_2	S_3	S_4	S_5	S_6	S_7	S_8
β_0	478	32	9	2	1	1	1	1
β_1	0	115	18	4	1	1	1	1

These numbers are consistent with the sample coming from some region with the homotopy type of a circle.

During an inclusion $S_i \hookrightarrow S_j$ holes can




During an inclusion $S_i \hookrightarrow S_j$ holes can



β_n^{ij} = number of n -dimensional holes in S_i that persist to S_j

During an inclusion $S_i \hookrightarrow S_j$ holes can



β_n^{ij} = number of n -dimensional holes in S_i that persist to S_j

β_0^{ij} = number of path components in S_i that persist to S_j


Bar codes


The matrix (β_n^{ij}) can be represented by a graph with horizontal edges and vertices arranged in columns.

The i th column has $\beta_n^{ii} = \beta_n(S_i)$ vertices.

There are β_n^{ij} paths from the i th column to the j th column.








β_0 bar codes



could be enhanced to dendrograms (or phylogenetic trees)




From data to spaces

Given a set $S \subset \mathbb{E}^n$ and number $\epsilon > 0$ consider

$$X_\epsilon^S = \bigcup_{x \in S} B(x, \epsilon)$$

with $B(x, \epsilon)$ the ball of radius ϵ centred on x .




From data to spaces


Given a set $S \subset \mathbb{E}^n$ and number $\epsilon > 0$ consider

$$X_\epsilon^S = \bigcup_{x \in S} B(x, \epsilon)$$


with $B(x, \epsilon)$ the ball of radius ϵ centred on x .



Any $0 < \epsilon_1 < \epsilon_2 < \epsilon_3 < \dots$ yields inclusions $X_{\epsilon_1}^S \subset X_{\epsilon_2}^S \subset X_{\epsilon_3}^S \subset \dots$




simeq



vertices \leftrightarrow coloured regions

edges \leftrightarrow intersections



simeq





edges \leftrightarrow (non-contractible) intersections

Cech complex NX_ϵ^S

NX_ϵ^S is the simplicial complex with vertex set S and one simplex for each $\sigma \subseteq S$ satisfying



$$\bigcap_{x \in \sigma} B(x, \epsilon) \neq \emptyset.$$



Cech complex NX_ϵ^S

NX_ϵ^S is the simplicial complex with vertex set S and one simplex for each $\sigma \subseteq S$ satisfying

$$\bigcap_{x \in \sigma} B(x, \epsilon) \neq \emptyset.$$




Theorem: $X_\epsilon^S \simeq NX_\epsilon^S$

Vietoris-Rips complex R_ϵ^S

R_ϵ^S is the simplicial complex with vertex set S and one simplex for each $\sigma \subseteq S$ satisfying


$$\text{distance}(x, y) \leq \epsilon \text{ for each pair } x, y \in \sigma.$$



Vietoris-Rips complex R_ϵ^S

R_ϵ^S is the simplicial complex with vertex set S and one simplex for each $\sigma \subseteq S$ satisfying


$$\text{distance}(x, y) \leq \epsilon \text{ for each pair } x, y \in \sigma.$$



Theorem: $R_\epsilon^S \subseteq NX_{\epsilon\sqrt{2}}^S \subseteq R_{\epsilon\sqrt{2}}^S$

A *simplicial complex* K is a collection of subsets of some (ordered) set S satisfying:

- ▶ $\{x\} \in K$ for all $x \in S$,
- ▶ if $\sigma' \subset \sigma \in K$ then $\sigma' \in K$.



Computing Betti numbers

The chain complex $C_*(K)$

$$\longrightarrow C_n(K) \xrightarrow{\partial_n} C_{n-1}(K) \longrightarrow \dots \xrightarrow{\partial_1} C_0(K) \xrightarrow{\partial_0} 0$$

- ▶ $C_n(K)$ = vector space with basis $\{\sigma \in K : |\sigma| = n+1\}$

▶

$$\partial_n(\{x_1, \dots, x_{n+1}\}) = \sum_{i=1}^{n+1} (-1)^i \{x_1, \dots, \hat{x}_i, \dots, x_{n+1}\}$$

Computing Betti numbers

The chain complex $C_*(K)$

$$\longrightarrow C_n(K) \xrightarrow{\partial_n} C_{n-1}(K) \longrightarrow \dots \xrightarrow{\partial_1} C_0(K) \xrightarrow{\partial_0} 0$$

- ▶ $C_n(K)$ = vector space with basis $\{\sigma \in K : |\sigma| = n+1\}$



$$\partial_n(\{x_1, \dots, x_{n+1}\}) = \sum_{i=1}^{n+1} (-1)^i \{x_1, \dots, \hat{x}_i, \dots, x_{n+1}\}$$

determines the *homology*

$$H_n(K) = \frac{\ker(\partial_n)}{\text{image}(\partial_{n+1})}$$

Computing Betti numbers

The chain complex $C_*(K)$

$$\longrightarrow C_n(K) \xrightarrow{\partial_n} C_{n-1}(K) \longrightarrow \dots \xrightarrow{\partial_1} C_0(K) \xrightarrow{\partial_0} 0$$

- ▶ $C_n(K)$ = vector space with basis $\{\sigma \in K : |\sigma| = n+1\}$



$$\partial_n(\{x_1, \dots, x_{n+1}\}) = \sum_{i=1}^{n+1} (-1)^i \{x_1, \dots, \hat{x}_i, \dots, x_{n+1}\}$$

determines the *homology*

$$H_n(K) = \frac{\ker(\partial_n)}{\text{image}(\partial_{n+1})}$$

and *Betti numbers*

$$\beta_n(K) = \dim(H_n(K)).$$

Filtrations

$$K_1 \subseteq K_2 \subseteq K_3 \subseteq \cdots \subseteq K_N$$

$$C_*(K_1) \subseteq C_*(K_2) \subseteq C_*(K_3) \subseteq \cdots \subseteq C_*(K_N)$$

determine *persistent homology*

$$H_n^{ij}(K) = \frac{\ker(\partial_n^i)}{\text{image}(\partial_{n+1}^j) \cap C_n(K_i)}$$

Filtrations

$$K_1 \subseteq K_2 \subseteq K_3 \subseteq \cdots \subseteq K_N$$

$$C_*(K_1) \subseteq C_*(K_2) \subseteq C_*(K_3) \subseteq \cdots \subseteq C_*(K_N)$$

determine *persistent homology*

$$H_n^{ij}(K) = \frac{\ker(\partial_n^i)}{\text{image}(\partial_{n+1}^j) \cap C_n(K_i)}$$

and *persistent Betti numbers*

$$\beta_n^{ij} = \dim(H_n^{ij}(K)).$$

Filtrations

$$K_1 \subseteq K_2 \subseteq K_3 \subseteq \cdots \subseteq K_N$$

$$C_*(K_1) \subseteq C_*(K_2) \subseteq C_*(K_3) \subseteq \cdots \subseteq C_*(K_N)$$

determine *persistent homology*

$$H_n^{ij}(K) = \frac{\ker(\partial_n^i)}{\text{image}(\partial_{n+1}^j) \cap C_n(K_i)}$$

and *persistent Betti numbers*

$$\beta_n^{ij} = \dim(H_n^{ij}(K)).$$

Theorem: All β_n^{ij} can be determined from semi-echelon forms of the two matrices ∂_n^N and ∂_{n+1}^N .

Example

$$\partial_n^4 = \left(\begin{array}{cccc|ccc|cc|ccc} x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ \hline & & & & x & x & x & & x & x & & x & x & x \\ & & & & x & x & x & & x & x & & x & x & x \\ & & & & x & x & x & & x & x & & x & x & x \\ \hline & & & & & & & & x & x & & x & x & x \\ & & & & & & & & x & x & & x & x & x \\ & & & & & & & & x & x & & x & x & x \\ \hline & & & & & & & & & & & x & x & x \\ & & & & & & & & & & & x & x & x \end{array} \right)$$

Example

$$\partial_n^4 = \left(\begin{array}{cccc|ccc|cc|ccc} x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ x & x & x & x & x & x & x & & x & x & & x & x & x \\ \hline & & & & x & x & x & & x & x & & x & x & x \\ & & & & x & x & x & & x & x & & x & x & x \\ & & & & x & x & x & & x & x & & x & x & x \\ \hline & & & & & & & & x & x & & x & x & x \\ & & & & & & & & x & x & & x & x & x \\ & & & & & & & & x & x & & x & x & x \\ \hline & & & & & & & & & & & x & x & x \\ & & & & & & & & & & & x & x & x \end{array} \right)$$

$$\dim(C_n(K_4)) = 12, \quad \dim(C_{n-1}(K_4)) = 14, \quad \dim(C_n(K_3)) = 9, \quad \dots$$

Column reduce

$$\partial_n^4 = \left(\begin{array}{ccc|ccc|ccc} x & x & & x & x & x & x & x & x \\ & x & & x & x & x & x & x & x \\ & & x & & x & x & x & x & x \\ & & & x & x & & x & x & x \\ & & & & x & x & x & x & x \\ & & & & & x & x & x & x \\ & & & & & & x & x & x \\ \hline & & & & x & x & x & x & x \\ & & & & & x & x & x & x \\ & & & & & & x & x & x \\ \hline & & & & & & x & x & x \\ & & & & & & & x & x \\ & & & & & & & & x \\ \hline & & & & & & & x & x \\ & & & & & & & & x \end{array} \right)$$

Column reduce

$$\partial_n^4 = \left(\begin{array}{cc|ccc} x & x & & & & & & & \\ & x & & x & x & x & x & x & x & x & x \\ & x & & & x & x & x & x & x & & x \\ & x & & & x & x & x & x & & & x \\ & x & & & x & x & x & x & & & x \\ & & & & x & x & x & x & & & x \\ \hline & & & x & x & & x & x & & & x \\ & & & & x & & x & x & & & x \\ & & & & & & x & x & & & x \\ \hline & & & & & & x & & & & x \\ & & & & & & x & & & & x \\ & & & & & & x & & & & x \\ \hline & & & & & & & & & x & \\ & & & & & & & & & x & \end{array} \right)$$

Column reduce

$$\partial_n^4 = \left(\begin{array}{cc|ccc} x & x & x & x & x & x & x & x \\ & x & x & x & x & x & x & x \\ & x & & x & x & x & x & x \\ & x & & x & x & x & x & x \\ & x & & x & x & x & x & x \\ & & & x & x & x & x & x \\ \hline & & & x & x & x & x & x \\ & & & & x & x & x & x \\ & & & & x & x & x & x \\ \hline & & & & & x & x & x \\ & & & & & x & x & x \\ & & & & & x & x & x \\ \hline & & & & & & x & x \\ & & & & & & & x \end{array} \right)$$

$$\dim(\ker(\partial_n^4)) = 3, \quad \dim(\text{image}(\partial_n^4) \cap C_{n-1}(K_2)) = 7, \quad \dots$$

Illustration: Natural Images

van Hateren & van der Schaaf: 4167 digital photos of random outdoor scenes.

Illustration: Natural Images

van Hateren & van der Schaaf: 4167 digital photos of random outdoor scenes.

Mumford, Lee, Pedersen: 8 000 000 random high-contrast 3×3 patches in \mathbb{E}^9 .





Illustration: Natural Images

van Hateren & van der Schaaf: 4167 digital photos of random outdoor scenes.

Mumford, Lee, Pedersen: 8 000 000 random high-contrast 3×3 patches in \mathbb{E}^9 .



By normalizing with respect to brightness and contrast the patches are projected onto a set of points \mathcal{M} in a topological seven-sphere $S^7 \subset \mathbb{E}^8$.

Carlsson, Ishkanov, de Silva, Zomorodian: consider high-density subsets $\mathcal{M}(k) \subset \mathcal{M}$.

Carlsson, Ishkanov, de Silva, Zomorodian: consider high-density subsets $\mathcal{M}(k) \subset \mathcal{M}$.

Codensity function: $\delta_k(x)$ is the euclidean distance from $x \in \mathcal{M}$ to its k th nearest neighbour in \mathcal{M} .


Carlsson, Ishkanov, de Silva, Zomorodian: consider high-density subsets $\mathcal{M}(k) \subset \mathcal{M}$.

Codensity function: $\delta_k(x)$ is the euclidean distance from $x \in \mathcal{M}$ to its k th nearest neighbour in \mathcal{M} .

$\mathcal{M}(k)$ contains the 25% of patches with least $\delta_k(x)$.


The β_1 bar code for a random sampling of $\mathcal{M}(300)$ yields a single persistent homology class.

The β_1 bar code for a random sampling of $\mathcal{M}(300)$ yields a single persistent homology class.




The β_1 bar code for a random sampling of $\mathcal{M}(15)$ yields five persistent homology classes.

The β_1 bar code for a random sampling of $\mathcal{M}(15)$ yields five persistent homology classes.




The β_1 bar code for a random sampling of $\mathcal{M}(15)$ yields five persistent homology classes.



The β_2 bar codes are less robust, but seem to indicate a single homology class with mod 2 coefficients.


The β_2 bar codes are less robust, but seem to indicate a single homology class with mod 2 coefficients.

$$H_2(\text{Klein bottle}, \mathbb{Z}_2) = \mathbb{Z}_2$$



The β_2 bar codes are less robust, but seem to indicate a single homology class with mod 2 coefficients.

$$H_2(\text{Klein bottle}, \mathbb{Z}_2) = \mathbb{Z}_2$$



Better formulations of persistence

$$H_*(K_1) \xrightarrow{t_1} H_*(K_2) \xrightarrow{t_2} \cdots \xrightarrow{t_{N-1}} H_*(K_N)$$

Better formulations of persistence

$$H_*(K_1) \xrightarrow{t_1} H_*(K_2) \xrightarrow{t_2} \cdots \xrightarrow{t_{N-1}} H_*(K_N)$$

can be encoded as a (graded) $\mathbb{F}[t]$ -module :

$$V = \bigoplus_{i=1}^N H_*(K_i)$$

Better formulations of persistence

$$H_*(K_1) \xrightarrow{t_1} H_*(K_2) \xrightarrow{t_2} \cdots \xrightarrow{t_{N-1}} H_*(K_N)$$

can be encoded as a (graded) $\mathbb{F}[t]$ -module :

$$V = \bigoplus_{i=1}^N H_*(K_i)$$

with

$$t \cdot (a_1, \dots, a_N) = (0, t_1(a_1), \dots, t_{N-1}(a_{N-1})).$$

Since $\mathbb{F}[t]$ is a euclidean domain:

Theorem

$$V \cong \bigoplus_i t^{a_i} \cdot \mathbb{F}[t] \oplus \left(\bigoplus_j t^{b_j} \cdot (\mathbb{F}[t]/(t^{c_j} \cdot \mathbb{F}[t])) \right).$$

Since $\mathbb{F}[t]$ is a euclidean domain:

Theorem

$$V \cong \bigoplus_i t^{a_i} \cdot \mathbb{F}[t] \oplus \left(\bigoplus_j t^{b_j} \cdot (\mathbb{F}[t]/(t^{c_j} \cdot \mathbb{F}[t])) \right).$$

free summands: homology generators born at time a_i which live forever.

Since $\mathbb{F}[t]$ is a euclidean domain:

Theorem

$$V \cong \bigoplus_i t^{a_i} \cdot \mathbb{F}[t] \oplus \left(\bigoplus_j t^{b_j} \cdot (\mathbb{F}[t]/(t^{c_j} \cdot \mathbb{F}[t])) \right).$$

free summands: homology generators born at time a_i which live forever.

torsion summands: homology generators born at time b_j which die at time $b_j + c_j$.

Since $\mathbb{F}[t]$ is a euclidean domain:

Theorem

$$V \cong \bigoplus_i t^{a_i} \cdot \mathbb{F}[t] \oplus \left(\bigoplus_j t^{b_j} \cdot (\mathbb{F}[t]/(t^{c_j} \cdot \mathbb{F}[t])) \right).$$

free summands: homology generators born at time a_i which live forever.

torsion summands: homology generators born at time b_j which die at time $b_j + c_j$.

Remark: $\mathbb{Z}[t]$ and $\mathbb{F}[s, t]$ are not PIDs.

Quivers

A persistence module V is a representation of the quiver

$$1 \longrightarrow 2 \longrightarrow \dots \longrightarrow N$$

- vector space $V(x)$ to each object x
- homomorphism $V(a)$ to each arrow a .

Quivers

A persistence module V is a representation of the quiver

$$1 \longrightarrow 2 \longrightarrow \dots \longrightarrow N$$

- vector space $V(x)$ to each object x
- homomorphism $V(a)$ to each arrow a .

Direct sums

Two representations V, W have sum $V \oplus W$:

- $(V \oplus W)(x) = V(x) \oplus W(x)$
- $(V \oplus W)(a) =$ direct sum of linear maps $V(a), W(a)$.

Quivers

A persistence module V is a representation of the quiver

$$1 \longrightarrow 2 \longrightarrow \dots \longrightarrow N$$

- vector space $V(x)$ to each object x
- homomorphism $V(a)$ to each arrow a .

Direct sums

Two representations V, W have sum $V \oplus W$:

- $(V \oplus W)(x) = V(x) \oplus W(x)$
- $(V \oplus W)(a) =$ direct sum of linear maps $V(a), W(a)$.

Decomposables

A sum $V \oplus W$ of two non-trivial representations is *decomposable*.

Quivers

A persistence module V is a representation of the quiver

$$1 \longrightarrow 2 \longrightarrow \dots \longrightarrow N$$

- vector space $V(x)$ to each object x
- homomorphism $V(a)$ to each arrow a .

Direct sums

Two representations V, W have sum $V \oplus W$:

- $(V \oplus W)(x) = V(x) \oplus W(x)$
- $(V \oplus W)(a) =$ direct sum of linear maps $V(a), W(a)$.

Decomposables

A sum $V \oplus W$ of two non-trivial representations is *decomposable*.

Finite type

A quiver has *finite type* if it has only finitely many isomorphism types of indecomposable representations.

Gabriel's theorem

A connected quiver is of finite type if and only if its underlying graph is of type A_n , D_n , E_6 , E_7 , E_8 .

Gabriel's theorem

A connected quiver is of finite type if and only if its underlying graph is of type A_n , D_n , E_6 , E_7 , E_8 .

Boot strapping

Use bar codes for small sub samples S_1, S_2, \dots of a large sample $S \subset M$ to infer about M .


Gabriel's theorem

A connected quiver is of finite type if and only if its underlying graph is of type A_n , D_n , E_6 , E_7 , E_8 .

Boot strapping

Use bar codes for small sub samples S_1, S_2, \dots of a large sample $S \subset M$ to infer about M .

Difficulty: If bar codes for S_i suggest $\beta_1 = 1$ the situation could be M or M' .



Carlsson, de Silva: Consider inclusions

$$S_1 \rightarrow S_1 \cup S_2 \leftarrow S_2 \rightarrow S_2 \cup S_3 \leftarrow \dots \leftarrow S_N$$

and persistence in the *ziz-zag diagram*

$$H_n(R_\epsilon^{S_1}) \rightarrow H_n(R_\epsilon^{S_1 \cup S_2}) \leftarrow H_n(R_\epsilon^{S_2}) \rightarrow H_n(R_\epsilon^{S_2 \cup S_3}) \leftarrow \dots \leftarrow H_n(R_\epsilon^{S_N})$$

Carlsson, de Silva: Consider inclusions

$$S_1 \rightarrow S_1 \cup S_2 \leftarrow S_2 \rightarrow S_2 \cup S_3 \leftarrow \dots \leftarrow S_N$$

and persistence in the *ziz-zag diagram*

$$H_n(R_\epsilon^{S_1}) \rightarrow H_n(R_\epsilon^{S_1 \cup S_2}) \leftarrow H_n(R_\epsilon^{S_2}) \rightarrow H_n(R_\epsilon^{S_2 \cup S_3}) \leftarrow \dots \leftarrow H_n(R_\epsilon^{S_N})$$

Gabriel's theorem: The quiver

$$1 \rightarrow 2 \leftarrow 3 \rightarrow 4 \leftarrow \dots \leftarrow N$$

has indecomposable representations $\mathbb{I}(i, j)$ for $1 \leq i < j \leq N$:

$$\mathbb{I}(1, 2) : \mathbb{F} \xrightarrow{1} \mathbb{F} \leftarrow 0 \rightarrow 0 \leftarrow \dots \leftarrow 0$$

$$\mathbb{I}(1, 3) : \mathbb{F} \xrightarrow{1} \mathbb{F} \xleftarrow{1} \mathbb{F} \rightarrow 0 \leftarrow \dots \leftarrow 0$$


Other applications:

Medical images $f: \mathbb{M} \rightarrow \mathbb{R}$ could be analyzed using bar codes.

Other applications:

Medical images $f: \mathbb{M} \rightarrow \mathbb{R}$ could be analyzed using bar codes.

Consider a torus \mathbb{M} , height function f




and filtration

$$\mathbb{M}_r = f^{-1}([0, r]).$$

Other applications:

Medical images $f: \mathbb{M} \rightarrow \mathbb{R}$ could be analyzed using bar codes.

Consider a torus \mathbb{M} , height function f



and filtration

$$\mathbb{M}_r = f^{-1}([0, r]).$$