

We consider the notions of *finite* and *infinite* sets, and the *cardinality* of a set.

Reasonable goals for this section are to become familiar with these ideas and to practice interpreting descriptions of sets that are presented in terse mathematical notation (this means, amongst other things, distinguishing between different kinds of brackets: $\{$, $[$, $($, etc.).

Definition

A set is *finite* if it is possible to list its distinct elements one by one, and this list comes to an end.

A set is *infinite* if any attempt at listing its distinct elements continues indefinitely.

Example

$\{1, 2, 3, 4, 5\}$ is a *finite* set - its only elements are the integers 1, 2, 3, 4, 5, there are five of them.

Definition

A set is *finite* if it is possible to list its distinct elements one by one, and this list comes to an end.

A set is *infinite* if any attempt at listing its distinct elements continues indefinitely.

Example

The interval $[1, 3]$ is an *infinite* set - it consists of *all* the real numbers that are at least equal to 1 and at most equal to 3.

$$[1, 3] := \{x \in \mathbb{R} : 1 \leq x \leq 3\}.$$

Note: The symbol “:=” here means this is a statement of the *definition* of $[1, 3]$.

Definition

A set is *finite* if it is possible to list its distinct elements one by one, and this list comes to an end.

A set is *infinite* if any attempt at listing its distinct elements continues indefinitely.

Example

\mathbb{Z} and \mathbb{Q} are *infinite* sets.

Definition

A set is *finite* if it is possible to list its distinct elements one by one, and this list comes to an end.

A set is *infinite* if any attempt at listing its distinct elements continues indefinitely.

Example

The set of real solutions of the equation

$$x^5 + 2x^4 - x^2 + x + 17 = 0$$

is a *finite* set. We don't know how many elements it has, but it has at most five, since each one corresponds to a factor of degree 1 of this polynomial of degree 5.

Definition

A set is *finite* if it is possible to list its distinct elements one by one, and this list comes to an end.

A set is *infinite* if any attempt at listing its distinct elements continues indefinitely.

Example

The set of prime numbers is *infinite*.

A pair of *twin primes* is a pair of primes that differ by 2: e.g. 3 and 5, 11 and 13, 59 and 61. It is not known whether the set of pairs of twin primes is finite or infinite.

Definition

The *cardinality* of a finite set S , denoted $|S|$, is the number of elements in S .

Example

- If $S = \{5, 7, 8\}$ then $|S| = 3$.

Definition

The *cardinality* of a finite set S , denoted $|S|$, is the number of elements in S .

Example

- If $S = \{5, 7, 8\}$ then $|S| = 3$.
- $|\{4, 10, \pi\}| = 3$

Definition

The *cardinality* of a finite set S , denoted $|S|$, is the number of elements in S .

Example

- If $S = \{5, 7, 8\}$ then $|S| = 3$.
- $|\{4, 10, \pi\}| = 3$
- $|\{x \in \mathbb{Z} : \pi < x < 3\pi\}| = 6$.
 Note: $\{x \in \mathbb{Z} : \pi < x < 3\pi\} = \{4, 5, 6, 7, 8, 9\}$.

Definition

The *cardinality* of a finite set S , denoted $|S|$, is the number of elements in S .

Example

- If $S = \{5, 7, 8\}$ then $|S| = 3$.
- $|\{4, 10, \pi\}| = 3$
- $|\{x \in \mathbb{Z} : \pi < x < 3\pi\}| = 6$.
 Note: $\{x \in \mathbb{Z} : \pi < x < 3\pi\} = \{4, 5, 6, 7, 8, 9\}$.
- The cardinality of \mathbb{Q} is infinite.

Remarks

Remarks

- The notation " $|\cdot|$ " is severely overused in mathematics. If x is a real number, $|x|$ means the absolute value of x . If S is a set, $|S|$ means the cardinality of S . If A is a matrix $|A|$ means the determinant of A . It is supposed to be clear from the context what is meant.

- The notation " $|\cdot|$ " is severely overused in mathematics. If x is a real number, $|x|$ means the absolute value of x . If S is a set, $|S|$ means the cardinality of S . If A is a matrix $|A|$ means the determinant of A . It is supposed to be clear from the context what is meant.
- Defining the concept of cardinality for infinite sets is trickier, since you can't say how many elements they have. We will be able to say though what it means for two infinite sets to have the same (or different) cardinalities.

Example

In a hotel, keys for all the guest rooms are kept on hooks behind the reception desk. If a room is occupied, the key is missing from its hook because the guests have it. If the receptionist wants to know how many rooms are occupied, s/he doesn't have to visit all the rooms to check - s/he can just count the number of hooks whose keys are missing.

In this example, the occupied rooms are in **one-to-one correspondence** with the empty hooks. This means that each occupied room corresponds to **one and only one** empty hook, and each empty hook corresponds to **one and only one** occupied room. So the number of empty hooks is the same as the number of occupied rooms and we can count one by counting the other.

Definition

Suppose that A and B are sets. Then a **one-to-one correspondence** or a **bijective correspondence** between A and B is a pairing of each element of A with an element of B , in such a way that every element of B is matched to exactly one element of A .

Definition

Suppose that A and B are sets. A function $f: A \rightarrow B$ is called a **bijection** if

- Whenever a_1 and a_2 are different elements of A , $f(a_1)$ and $f(a_2)$ are different elements of B .
- Every element b of B is the image of some element a of A .

Cardinality and bijective correspondence

If a bijective correspondence exists between two finite sets, they have the same cardinality. Sometimes, in order to determine the cardinality of a set, it is easiest to determine the cardinality of another set with which we know it is in bijective correspondence.

Example

How many integers between 1 and 1000 are perfect squares?

Solution: The list of perfect squares in our range begins as follows

$$1, 4, 9, 16, \dots$$

One way of solving the problem would be to keep writing out successive terms of this sequence until we hit one that exceeds 1000, and then delete that one and count the terms that we have. This is actually more work than we are asked to do, since we are not asked for the list of squares but just the number of them.

Cardinality and bijective correspondence

If a bijective correspondence exists between two finite sets, they have the same cardinality. Sometimes, in order to determine the cardinality of a set, it is easiest to determine the cardinality of another set with which we know it is in bijective correspondence.

Example

How many integers between 1 and 1000 are perfect squares?

Solution:

Alternatively, we could notice that $(31)^2 = 961$ and $(32)^2 = 1024$. So the numbers $1^2, 2^2, \dots, (31)^2$ are all in the range 1 to 1000 and these are the only perfect squares in that range, the answer to our question is 31.

If a bijective correspondence exists between two finite sets, they have the same cardinality. Sometimes, in order to determine the cardinality of a set, it is easiest to determine the cardinality of another set with which we know it is in bijective correspondence.

Example

How many integers between 1 and 1000 are perfect squares?

Solution:

What is used here is the fact that the set of perfect squares in the range of interest is in bijective correspondence with the set $\{1, 2, 3, \dots, 31\}$ - it's not really the squares in the range 1 to 1000 that we are counting but the integers in the range 1 to 31.

This last example shows that it could be possible to know that there is a bijective correspondence between two finite sets, without knowing the cardinality of either of them.

Example

Show that the equations

$$x^3 + 2x + 4 = 0 \text{ and } x^3 + 3x^2 + 5x + 7 = 0$$

have the same number of real solutions.

$$x^3 + 2x + 4 = 0 \text{ and } x^3 + 3x^2 + 5x + 7 = 0$$

Learning outcomes for Section 2.2

Solution: One way of doing this is to demonstrate a bijective correspondence their sets of real equations. We can write

$$\begin{aligned} x^3 + 3x^2 + 5x + 7 &= (x^3 + 3x^2 + 3x + 1) + 2x + 6 \\ &= (x + 1)^3 + (2x + 2) + 4 \\ &= (x + 1)^3 + 2(x + 1) + 4. \end{aligned}$$

This means that a real number a is a solution of the second equation if and only if

$$(a + 1)^3 + 2(a + 1) + 4 = 0$$

i.e. if and only if $a + 1$ is a solution of the first equation.

The correspondence $a \leftrightarrow a + 1$ is a bijective correspondence between the solution sets of the two equations. So they have the same number of real solutions.

Note: This number is at least 1 and at most 3. Why?

After studying this section you should be able to

- Explain what is meant by the cardinality of a set;
- Read and interpret descriptions of different subsets of \mathbb{R} presented using different standard notations. Decide what the elements of these sets are and whether the sets are finite or infinite;
- Explain what is meant by a bijective correspondence and give examples to support your explanation.